

SYSTEM DERO: ROZWÓJ TECHNIK OPERACYJNEGO LECZENIA KREGOSŁUPA

A. Skwarcz, P. Majcher, M. Fatyga, S. Zaborek¹

Stabilization of the fractured spine, spondylolisthesis and Idiopathic Scoliosis with Transpedicular DERO System

*(Zastosowanie transpedikularnej stabilizacji
systemem DERO w skoliozach idiopatycznych,
urazach kręgosłupa i kręgozmykach)*

Summary. Basing on the clinical material the autors show the results of spine deformities treatment with transpedicular

screws. The use of transpedicular screws improved the deformities correction and spine stabilization.

List of figures:

- Fig.1. A girl K.D.No.820609. Juvenile Idiopathic right side thoracic and left side lumbar scoliosis, King II.
A. Preoperative A-P X-rays 62°/54°.
B. Postoperative A-P X-ray with the DERO System 23°/21°.
C. Postoperative lateral X-ray, transpedicular screws in L₁, L₂, L₃.
- Fig.2. A girl W.A.No.810818. Juvenile idiopathic thoracic right side scoliosis, King III.
A. Preoperative A-P X-ray 64°.
B. Postoperative A-P X-ray with the DERO System 15°.
C. Postoperative lateral X-ray transpedicular screws in L₁, L₂.
- Fig.3. Patient K.B.No.480531. L1 vertebral body compression fracture with kyphotic deformity of the spinal column and partly damaged spinal cord and roots. L1 laminectomy with the spinal cord decompression, correction of the kyphotic deformity and stabilization with DERO System. Spondylodesis with autogenous graft.
A. Sagittal reconstruction the fracture is seen.
B, C. Postoperative A-P and lateral X-ray.
- Fig.4. Pacient W.J.No.270104. Spondylolisthesis L₄-L₅. Reposition and stabilization with the DERO System.
A, B. Preoperative A-P and lateral X-ray.
C, D. Postoperative A-P and lateral X-ray.

Authors:

¹) Skwarcz Andrzej, Majcher Piotr, Fatyga Marek, Zaborek Sławomir. Katedra i Klinika Rehabilitacji Akademii Medycznej w Lublinie. Kierownik: Prof. dr hab. med. Andrzej Skwarcz

Corespondence and reprint request to:

S k w a r c z A n d r z e j

Katedra i Klinika Rehabilitacji Akademii Medycznej w Lublinie
Państwowy Szpital Kliniczny Nr 4
ul. Jacewskiego 8
20-950 LUBLIN
Poland